

ON THE RATIONALITY OF $SU(r, d)$

DAVID C. BUTLER

INTRODUCTION

Let C be a curve of genus g over the complex numbers. For $0 < d < r$, let $SU = SU(r, r(g-1) + d)$ be the moduli space of bundles with rank r and fixed determinant of $\deg r(g-1) + d$ over C . This paper shows that if d divides $(r \pm 1)$, then SU is rational (no matter what the value of g). In fact if the prime divisors of $\delta > 0$ are divisors of r , and d divides $r - \delta > 0$ then SU is rational for any genus. We conjecture that this holds for $r + \delta$ if $0 < \delta < r$.

Research on the rationality of SU began shortly after the construction of the Moduli space. In 1964, Tyurin [9] proved the case of rank 2 bundles of odd degree. He went on to claim rationality for all ranks and degrees [10], but the paper contained fatal errors. Newstead corrected some errors, and in his unpublished thesis, he proved the case $d = 1$ for any r . He then went on to claim a proof for $(r, d) = 1$ [6]. That proof contained a trivial but fatal error. He concluded in his correction [6] that for any pair (r, d) which are coprime the theorem holds for infinitely many genera. Which suggests, but does not prove, that the theorem is true. Ballico then noted that Newstead's argument proved SU was stably rational [1], where X is stably rational with level k if for all $n \geq k$, $X \times \mathbb{P}^n$ is rational. Unfortunately, Beauville, Colliot-Thélène, Sansuc and Swinnerton-Dyer proved stable rationality does not imply rationality [2]. However, Ballico noted that stable rationality with small level and Newstead's arguments would yield strong results. Aware of this, Boden and Yokogawa [3], used parabolic bundles to prove $SU(r, d)$ is stably rational

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

of level $\leq r - 1$. This, and the argument of Newstead proved that if $(r, d) = 1$, then SU is rational provided either $(g, d) = 1$ or $(g, r - d) = 1$. And this result generalizes Newstead's results.

Now our results continue the theme of widening the number of genera for which rationality is known. In particular, we find infinitely many pairs (r, d) with $(r, d) = 1$ where $SU(r, d)$ is rational.

Main Theorem. *Let C be a smooth projective irreducible curve over $\mathbb{C}$. Then $SU(r, d')$ is rational if $d' \equiv \pm d \pmod{r}$, $(r, d) = 1$ and if*

A) *d divides $(r \pm 1)$, or if*

B) *d divides $(r - \delta)$, where for all primes p such that $p|\delta$, $p|r$, and $0 < \delta < r$.*

Along the way we wind up proving a second, seemingly unrelated result:

Second Theorem. *Let C be as above, and let $\pi : \tilde{C} \rightarrow C$ be a cyclic etale cover of deg e . For a general stable bundle E with $\mu(\pi^*E) > g(\tilde{C})$, $\pi^*(E)$ is generated by global sections.*

Before we outline the proof, it should be noted that $SU(r, d) \cong SU(r, rk \pm d)$ for any integer k by tensoring and/or dualizing.

Now we outline the proof giving a historic perspective. It begins with Newstead who observed that since a general bundle with rank r and $\deg r(g - 1) + d$ had $h^0 = d$, there is a unique subbundle $I = \bigoplus^d \mathcal{O}_C$ to a general E . This gives an exact sequence

$$0 \rightarrow I \rightarrow E \rightarrow F \rightarrow 0.$$

It seems likely, and turns out to be true that for general E , F is a stable bundle, and so a general E is given by an extension of a rank $r - d = n$, $\deg r(g - 1) + d$ vector bundle F by I . Now we know $SU = SU(r, r(g - 1) + d)$ has a map to $SU' = SU(r - d, r(g - 1) + d)$ whose fiber over a closed point $[F]$ is $\text{Ext}^1(F, I) \cong H^0(C, \omega_C \otimes F)^*$ modulo the action of $\text{Aut}(I)$,

and this turns out to be a Grassmannian. But even if $SU' = SU(r - d, r(g - 1) + d)$ is rational, SU need not be rational. Because unfortunately, if X maps to a rational variety with rational fibers it is not necessarily true that X is rational. But if $(r - d, r(g - 1) + d) = 1$, then SU' has a Poincare bundle $\mathcal{P}$ on $SU' \times C$. Tensoring by the pullback of ω_C then pushing down to SU' and taking d direct summands, we get a Zariski open set with a bundle parameterizing $\text{Ext}^1(F, I)$. Taking $\mathbb{G}$ the quotient by $\text{Aut}(I)$, we get a parameterization of an open set of SU by a Grassmannian bundle over SU' . Since it is a bundle, it is trivial (on a perhaps smaller set) and so SU is birational to $SU' \times \mathbb{G}$. If SU' is rational by inductive hypothesis then SU is rational. Unfortunately there is one problem. $r(g - 1) + d$ and $r - d$ need not be relatively prime. And if they are not, there is no Poincare bundle.

To see that the above happens consider the following example. Set $r = 5$, $g = 6$ and $d = 2$. $5(6 - 1) + 2 = 27$ and is not relatively prime to $(5 - 2) = 3$. The same thing happens for $d = 5 - 2 = 3$. Now consider another example. Set $r = 7$, $g = 6$ and $d = 2$. $7(6 - 1) + 2 = 37$ which is relatively prime to $(7 - 2)$. But $SU' = SU(5, 5(6 - 1) + 2)$ which is the previous example. We can try $d = 7 - 2 = 5$ but $7(6 - 1) + 5 = 40$ is not relatively prime to $7 - 5 = 2$. The whole inductive argument breaks down unless g is chosen carefully, or if $d = \pm 1$, in which case substituting $d = r - 1$ (By tensoring and perhaps dualizing) gives a rank 1 cokernel which is rational because it is a point and there is only one fiber.

Now Ballico notes that in the coprime-prime case we do not need the SU' to be rational, but for some product of a rational variety with SU' to be rational. If SU' is stably rational with level less than or equal to the dimension of the Grassmannian, then SU is rational. Unfortunately Ballico could not prove the moduli spaces were stably rational of small enough level; although he did prove them stably rational.

Boden and Yokogawa then proved rationality for most parabolic bundles. As a corollary they proved that bundles with $(\text{rank}, \text{deg}) = (r, d) = 1$ are stably rational with level $r - 1$. This implies SU is stable if $(r(g - 1), r - d) = 1$ or $(r(g - 1), d) = 1$. And this simplifies

to $(g, d) = 1$ or $(g, r - d) = 1$.

Now the author enters the picture. It would seem at first that Newstead's original argument had reached its logical conclusion. But it could be improved by changing its very beginning. Newstead begins with the trivial bundle I of rank d , because its embedding into a general E with $[E] \in SU$ is unique up to automorphisms of I . But there are other bundles with this property. Namely any $\deg 0$ twist of I or any direct sum of $\deg 0$ twists of I . We use the latter. Let $\{L_1, \dots, L_e\}$ be a set of mutually non-isomorphic $\deg 0$ line bundles and let U_i be a rank d vector space. Then set $V_i = U_i \otimes L_i$ and $W = \bigoplus^e V_i$. The bulk of our proof is to show that for any W with $ed < r$ and a general E the sequence

$$0 \rightarrow W \rightarrow E \rightarrow F \rightarrow 0$$

has a stable bundle F as cokernel. And when $\text{rank } W > \text{rank } E$ we have a surjection of W to E giving an exact sequence,

$$0 \rightarrow K_{E,W} \rightarrow W \rightarrow E \rightarrow 0.$$

For the plus case in part A, $K_{E,W}$ is a fixed line bundle we call A . And SU is isomorphic to $H^0(C, A^* \otimes W) / \text{Aut}(W)$, and this turns out to be a product of Grassmannians.

Now for part B (which implies the minus case in part A). If $ed = r - \delta$ where every prime factor of δ is a prime factor of r , then $(\delta, r(g-1) + d) = 1$ since no divisor of r divides d . That means $SU' = SU(\delta, r(g-1) + d)$ has a Poincare bundle. A general E is an extension of $F \in SU'$ by W . And since SU' has a Poincare bundle, it has a bundle whose fibers are $\text{Ext}^1(F_t, W) \cong \bigoplus^i H^0(C, \omega_C \otimes L_i^* \otimes F_t)^* \otimes U_i$. Taking an affine subset of SU' we can form a universal extension space. This gives us a map from the bundle over SU' to SU which is dominant and has fibers $\text{Aut}(W)$. We take the quotient by the automorphism group and get a birational map from SU' cross a product of Grassmannians to SU . Since the product of Grassmannians has dimension $\geq r - 1$, SU is rational.

It would be nice to have similar results for when $ed = r + \delta$ but they are held up because we do not know when $K_{E,W}$ is stable. In the final section we make a natural conjecture and offer some evidence. We will also prove the Second Theorem.

I thank Johns Hopkins University and the Tata Institute for Fundamental Research for their hospitality. I would like to thank Shokurov who shared his expertise in birational geometry. I would also like to thank Mehta, Nitsure, Ramadas Ramanan and Srinivas for answering questions so numerous that I do not know who answered what. But I do know they were answered.

§1 PRELIMINARIES

C is a smooth projective irreducible curve of genus g over $\mathbb{C}$

For a vector bundle E , $\mu(E) = \deg(E)/\text{rank}(E)$.

E is stable (semistable) if $\mu(S) < \mu(E)$ ($\leq$) for all proper subbundles. If E is not semistable, then it is called unstable. If E is not stable, then it is called nonstable. If E is semistable but not stable, then it is called strictly semistable.

As in the introduction $SU = SU(r, r(g-1) + d)$ and $U = U(r, r(g-1) + d)$, the moduli spaces of stable bundles of rank r and $\deg r(g-1) + d$ with fixed and unfixed determinants respectively. It is assumed that $0 < d < r$.

By abuse of notation we say $E \in SU$ (or U) if the isomorphism class of the vector bundle is in the moduli space.

We will use E to denote a fixed bundle and E_t to denote a general bundle in the moduli space.

L will denote a single linebundle of $\deg = 0$.

We will consider sets of mutually non-isomorphic line bundles of $\deg 0$: $\{L_1, L_2, \dots, L_e\}$. Our results will not depend on the the choice (except for the Second Theorem).

$U_i = \bigoplus^d k$ where k is the ground field.

$V_i = U_i \otimes L_i$.

$$W_k = V_1 \oplus \cdots \oplus V_k.$$

$$W = W_e.$$

We will use dimension counts to prove our results. These arguments are hampered because many bundles do not live on a moduli space, and even those that do may not have a universal or Poincare bundle. But we may (after Sundaram [8]) form very useful parameter spaces of bundles.

Definition. Let Cl be a class of vector bundles over C . A parameter space T of Cl is a variety along with a bundle $\mathcal{E}$ over $T \times C$ such that any E in Cl is isomorphic to the restriction E_t of $\mathcal{E}$ to some $t \times C$.

Remark 1. It is not assumed that $E_t \cong E_s$ implies $s = t$. And even if it does, parameter spaces do not generally have the universal properties of a moduli space.

Stable vector bundles of rank r and $\deg d$ where $(r, d) = 1$ form a fine moduli space. In particular, there is a Poincare bundle $\mathcal{P}$ over the moduli spaces SU and U . In the non-coprime case, the variety is quasi-projective and has no Poincare bundle (even locally). However U and SU can be covered by Zariski open sets $\{A_i\}$. The A_i in turn have etale covers $p_i : T_i \rightarrow A_i$. There is a Poincare bundle $\mathcal{P}_i$ over T_i such that the map given by the universal property of U and SU is p_i . This gives a parameter space of stable bundles with rank r and degree d where $(r, d) \neq 1$.

Sundaram constructs parameter spaces from the Jordan Holder filtration, and the Harder Narasimhan filtration. The strictly semistable bundles are parameterized by finitely many components with dimension $\leq \dim M - (r - 1)$. The unstable bundles have infinitely many components. So this case needs explaining. First of all, for any Bundle E with rank r and degree d , there is a unique filtration:

$$\sum E : 0 = \mathcal{E}_1 \subset \mathcal{E}_2 \cdots \subset \mathcal{E}_k = E,$$

where $E_i = \mathcal{E}_i / \mathcal{E}_{i-1}$ is semistable, and $\mu_i = \mu(E_i)$ is strictly decreasing. We say that E has Harder Narasimhan (or HN) type $\{(r_1, \mu_1), (r_2, \mu_2), \dots, (r_k, \mu_k)\}$; $r_1 + r_2 + \cdots + r_k = r$, where

$r_i = \text{rank}(E_i)$. We set $\mu_1 = \mu^{\max}$; it is the upper bound on μ of a subbundle. Similarly, we set $\mu_k = \mu^{\min}$; it is the lower bound on μ of a quotient bundle. If $\mu^{\max} < 0$ then $h^0 = 0$ as $\mathcal{O}_C$ is not a subbundle. And by Serre Duality, if $\mu^{\min}(E) > 2g - 2$, then $h^1 = 0$. Since the complex numbers are characteristic zero, $\mu^{\max}(E \otimes F) = \mu^{\max}(E) + \mu^{\max}(F)$. $\mu^{\min}(E \otimes F) = \mu^{\min}(E) + \mu^{\min}(F)$. Also $\mu^{\max}(E^*) = -\mu^{\min}(E)$ by dualizing the HN filtration. and likewise $\mu^{\min}(E^*) = -\mu^{\max}(E)$.

If r is fixed and $\mu^{\max}$ is bounded from above and $\mu^{\min}$ is bounded from below, there are only finitely many HN types possible. This is also the case when $\deg$ is bounded from above and $\mu^{\min}$ is bounded from below, or if $\mu^{\max}$ is bounded from above and $\deg$ is bounded from below. The advantage is that then there are only finitely many components of the parameter space of unstable bundles, and each has dimension $\leq \dim U - (r - 1)$ or $\leq \dim SU - (r - 1)$.

§2 SOME PROPOSITIONS

Proposition 1. *Let $0 < c \leq d$ and $c < r$ and L be a line bundle of $\deg = 0$. For a general $E_t \in U(r, r(g - 1) + d)$ with $d > 0$ and d not necessarily $< r$, there is an exact sequence:*

$$0 \rightarrow \bigoplus^c L \rightarrow E_t \rightarrow F_u \rightarrow 0$$

with F_u stable.

Remark 2. This does not mean that all embeddings of $\bigoplus^c L$ have a stable vector bundle cokernel. But a general subbundle of a general bundle such a cokernel. Of course if $c = d$ there is only one such subbundle for general E_t and the cokernel is unique.

Proof of Proposition 1. The first thing that can go wrong is that the cokernel is not a bundle. In some sense this is all that can go wrong outside of stability. If $c = 1$ this is obvious. If $c = 2$, the one dimensional family of (twisted) sections may span a subbundle of E_t of rank 1. But going back to $c = 1$, that means L factored. If that never happened,

there is no problem at $c = 2$ except maybe factoring through a subbundle of rank 2, and so on. So we can assume $\overset{c}{\oplus}L$ is a subbundle.

So now the idea is to count dimensions of the space of subbundles in various E_t that our trivial bundles could factor through and the dimensions of the space of subbundles of the form $\overset{c}{\oplus}L$. We know $\overset{c}{\oplus}L$ embeds in every E_t . We also know $h^0(C, E_t) = d$ for general t and there are $c(d - c)$ dimensions of embeddings. So the total dimension of subbundles of the form $\overset{c}{\oplus}L$ is

$$(*) \quad r^2(g - 1) + 1 + c(d - c),$$

which we will show is larger than the dimension of the space of bundles factored through.

So suppose $\overset{c}{\oplus}L$ factors through a bundle S and that we have a sequence;

$$0 \rightarrow S \rightarrow E_t \rightarrow Q \rightarrow 0$$

where Q is a vector bundle. S is a rank c bundle and an elementary transformation of $\overset{c}{\oplus}L$. So at first glance the dimension of possible bundles S is $c\delta$ with $\delta = \deg S$, since S must be an elementary transform of $\overset{c}{\oplus}L$. But there are $c^2 - 1$ projective automorphisms of $\overset{c}{\oplus}L$. So if $\delta \geq c$ the dimension is $c\delta - c^2 + 1$. If $\delta < c$ the dimension is 0 and there are at least $c^2 - c\delta - 1$ dimension of automorphism of S . The dimension of automorphisms must be subtracted from the dimension of extensions, so we shall abuse terminology and say the dimension of the space of bundles S with a map $\overset{c}{\oplus}L \rightarrow S$ is $c\delta - c^2 + 1$.

We know that

$$\deg Q = r(g - 1) + d - \delta = (r - c)(g - 1) + c(g - 1) + d - \delta.$$

And since $\mu^{\min}(Q) \geq \mu(E_t) > 0$ is bounded from below, only finitely many HN types are possible for Q . The dimension is bounded by the dimension for stable Q , that is

$(r-c)^2(g-1)+1$. Since Q is a quotient of the stable bundle E_t . $\mu^{\min}(Q) > \mu(E_t)$. Similarly $\mu^{\max}(S) < \mu(E_t)$ And hence $\mu^{\min}(S^*) > -\mu(E_t)$. All of which gives us $\mu^{\min}(\omega_C \otimes S^* \otimes Q) > 2g - 2$ And hence, $h^1(C, \omega_C \otimes S^* \otimes Q) = 0$.

Now we estimate and bound the dimension of the space of sections.

$$h^0(C, \omega_C \otimes S^* \otimes Q) = 2c(r-c)(g-1) - \delta(r-c) + c^2(g-1) + cd - c\delta.$$

Now we add the dimension of bundles S and Q then subtract 1 from Ext^1 in order to get the projective dimension. It all comes out to:

$$(**) \quad r^2(g-1) + 1 + cd - c^2 - \delta(r-c).$$

The upshot is that $*$ is bigger than $**$, so $\overset{c}{\oplus} L$ is a subbundle with a bundle cokernel.

To see that the general cokernel is stable, consider yet another dimension count. We count extensions of the form

$$0 \rightarrow \overset{c}{\oplus} L \rightarrow E_t \rightarrow F_u \rightarrow 0.$$

Let $F_u \in U$ be a parameter space of stable bundles of rank $r-c$ and $\deg r(g-1) + d$. It has dimension $(r-c)^2(g-1) + 1$. Calculating as above we see the dimension of extensions is $r^2(g-1) + c(d-c)$. A parameter space of nonstable bundles would have dimension at least $n-1$ lower; therefore, the general cokernel is stable. That proves Proposition 1. $\square$

Proposition 2. *Suppose that for a general bundle $E_t \in U$ there is a natural sequence*

$$0 \rightarrow W_k \rightarrow E_t \rightarrow F_u \rightarrow 0$$

where F_u is a stable vector bundle for general u . If $\overset{c}{\oplus} L$ (with $\deg L = 0$) embeds in a general E_t , $L \not\subseteq W_k$, and $dk + c < r$, then for (perhaps more general E_t) there is an exact sequence:

$$0 \rightarrow W_k \overset{c}{\oplus} L \rightarrow E_t \rightarrow F'_v \rightarrow 0$$

with F'_v a stable vector bundle and $\overset{c}{\oplus}L$ a general subbundle of E_t . Hence, by setting $c = d$ and considering only E_t with $h^0(C, E_t \otimes L_i^*) = d$, and with $ed < r$, we get for general t , a unique sequence:

$$0 \rightarrow W_e \rightarrow E_t \rightarrow F_u \rightarrow 0$$

with F_u a stable bundle for general t .

Proof of Proposition 2. First we start with $F_u = F$ fixed. Let $\overset{c}{\oplus}L$ be a subbundle of F . Then for a general subbundle Proposition 1 applies and the cokernel is stable. Now we have to pullback the subbundle of F to a subbundle of E_t .

For any given embedding $\overset{c}{\oplus}L \xrightarrow{\alpha} F$, there is a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & W_k & \longrightarrow & E_{t\alpha} & \longrightarrow & \overset{c}{\oplus}L \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \alpha \\ 0 & \longrightarrow & W_k & \longrightarrow & E_t & \longrightarrow & F \longrightarrow 0 \end{array}$$

$E_{t\alpha}$ is base extension of E_t over F by α . So α factors through E_t iff it factors through $E_{t\alpha}$, which means the top sequence splits. To interpret this, consider the surjective induced map:

$$H^0(C, \omega \otimes W_k^* \otimes F)^* \twoheadrightarrow H^0(C, \omega \otimes W_k^* \otimes (\overset{c}{\oplus}L))^*.$$

The top sequence splits if the element corresponding to the bottom sequence maps to 0. It is easily verified by a dimension count that there are extensions which cause the upper sequence to split, and many of them. For the kernel is the dimension of the bottom extensions of F to W_k minus the dimension of top extensions $\overset{c}{\oplus}L$ to W_k .

We can choose E_t so it has $h^1(C, E_t \otimes L_i^*) = 0$ for $1 \leq i \leq k$ and $h^1(C, E_t \otimes L^*) = 0$. This forces a choice of F but we can choose E_t so that F is stable. Also note, the general extension of F by W_k will be stable and have the above vanishings of h^1 . Now we count the dimension of pairs, $(\overset{c}{\oplus}L \subset E_t)$ denoting a specific subbundle $\overset{c}{\oplus}L$. It's dimension is the

dimension of the bottom extensions plus $c(d - c)$. Now this maps to the set of subbundles of F . So we subtract the dimension of the kernel, which means we subtract the dimension of the bottom extensions and add the number of top extensions. This gives us a total of $c(d - c) + kcd(g - 1)$.

Now we count the dimension of pairs $(\overset{c}{\oplus} L \subset F)$. Since a general E_t has $h^1(C, E_t \otimes L^*) = 0$, so does F have $h^1(C, F \otimes L^*) = 0$. Since $h^0(C, F \otimes L^*) = r(g - 1) + d - (r - dk)(g - 1)$, we get dimension of rank c subbundles given by, $cdk(g - 1) + c(d - c)$. So the map is dominant and a general bundle $\overset{c}{\oplus} L \subset E_t$ has a stable cokernel (provided our bundle is an extension of F).

The preceding argument has two faults. The first is that we never constructed the object whose dimensions we counted. The second is that it handles F_u one u at a time, treating the family of F_u as a set (with no algebraic structure).

For the first, let $V = \text{Ext}^1(F, W_k)$. Now there is a sequence over $V \times C$:

$$0 \rightarrow \mathcal{W}_k \rightarrow \mathcal{E} \rightarrow \mathcal{F} \rightarrow 0.$$

Given a point $v \in V$ the sequence restricts over $v \times C$ to:

$$0 \rightarrow W_k \rightarrow E_t \rightarrow F \rightarrow 0,$$

where the extension corresponds to $v \in V$. See [5] [7] and [4]. If we tensor by L^* and push down $\pi : V \times C \rightarrow V$, we get a sequence

$$0 \rightarrow \pi_* \mathcal{E}' \rightarrow \pi_* \mathcal{F}'.$$

When restricted to $v \times C$ we get:

$$0 \rightarrow H^0(C, E_t) \rightarrow H^0(C, F).$$

A rank c subspace in $H^0(C, E_t)$ corresponds to a unique subbundle $\overset{c}{\oplus} L \subset E_t$. And its image in $H^0(C, F)$ corresponds to the subbundle's image in F . So we take the Grassmannian

bundles and note that the second Grassmannian bundle maps from $V \times Gr$ to Gr . So a Grassmannian bundle over V dominates a Grassmannian.

The previous dimension count now holds. And our dimension count has shown a birational map between subbundles of general E_t and those of F . We have a general subbundle, $\overset{c}{\oplus} L \subset F$ has a stable cokernel by Proposition 1. And that implies a general subbundle $W_k \overset{c}{\oplus} L \subset E_t$ has a stable cokernel provided E_t has cokernel F corresponding to $W_k \subset E_t$.

We need to extend this result to a general cokernel F_u . So we need a general construction of a universal extension of two families of vector bundles. We have W_k which does not really vary, and F_u which does. So we have two families T and S . T is a point, and S is some cover of an open subset of SU the moduli of stable bundles of some fixed rank and degree. First we need $H^1(C, F_u \otimes L^*) = 0$. This is true for general F_u . Then we need to take an open set and an etale cover to get a Poincare bundle. We then take S a Zariski open affine subspace of the etale cover. This will still dominate. We now satisfy the criteria of [Lemma 2.4, 7], because an affine space has no higher cohomology. So by that lemma, there is a universal extension with parameter $\mathbb{P}(V)$ which is a projective bundle over U whose fiber at u is $H^1(C, \text{Hom}(F_u, W_k))$. It can be formed by taking the pullback of $\omega_C \otimes W^*$, tensoring by the Poincare bundle $\mathcal{P}$ and pushing down onto U and dualizing. Now we can take Grassmannian bundles and globalize the proof, showing that for a general subbundle $\overset{c}{\oplus} L \subset E_t$ of a general bundle E_t the cokernel is a stable bundle. $\square$

Proposition 3. *For a general bundle $E_t \in U$, a line bundle L_{k+1} of $\deg 0$ with $L_{k+1} \not\subset W$, a general $\overset{c}{\oplus} L_{k+1} \subset E_t$ with $c \leq d$, and $W_k \subset E_t$, the natural map*

$$W_k \overset{c}{\oplus} L_{k+1} \rightarrow E_t$$

is surjective provided $dk + c \geq r + 1$.

Proof of Proposition 3. To begin with we show that the map is at least generically surjective for a general E_t . So let i be the largest integer such that $di \leq r - 1$. Then for a general

E_t and a general $\overset{c}{\oplus} L_{i+1} \subset E_t$, with $c' = r - 1 - di$, we get a subbundle whose quotient is a line bundle, by Proposition 2. Now another subbundle $L_{i+1} \subset E_t$ which does not factor through $W_i \overset{c'}{\oplus} L$ maps to the line bundle quotient generically. We can choose another L_{i+2} , set $c' + 2 = c$, and the map $W_k \overset{c}{\oplus} L_{k+1}$ generically surjects onto E_t .

Now we have to show surjectivity for a general E_t . Once again, this will be done by dimension count. First we count the number of pairs $W_k \overset{c}{\oplus} L_{k+1} \subset E_t$. The dimension of the E_t is $r^2(g - 1) + 1 = (r^2 - 1)(g - 1) + g$. The dimension of the subbundle (for given E) is the dimension of the Grassmannian $(c, d) = cd - c^2$. So our final dimension is:

$$(***) \quad (r^2 - 1)(g - 1) + g + cd - c^2.$$

Now suppose the map in the proposition does not surject onto E_t . It then surjects onto some bundle Q and E_t is an elementary transformation of Q . So this means

$$\deg Q = \deg E_t - \delta = r(g - 1) + d - \delta$$

To calculate the dimension of the family of Q 's consider the sequence:

$$0 \rightarrow K \rightarrow W_k \overset{c}{\oplus} L_{k+1} \rightarrow Q \rightarrow 0,$$

where K is a line bundle, with $\deg K = r(g - 1) + d - \delta$. Assume for now $h^1(C, K^* L_i) = 0$ for all L_i and K . An informal dimension count gives:

$$(r + 1)(r - 1)(g - 1) + g + (r + 1)d - (r + 1)\delta.$$

This follows from there being $r + 1$ subline bundles in $W_k \overset{c}{\oplus} L_{k+1}$. All of them have $\deg = 0$, and K^* has $\deg = r(g - 1) + d - \delta$, and therefore, $h^0(C, K^* \otimes L_i) = (r - 1)(g - 1) + d - \delta$. The g is the dimension of the possible K . Now we need to account for the automorphisms which come to $kd^2 - c^2$. Now note that $(r + 1)d = (kd + c)d = (kd^2 + cd)$. And hence,

$(r+1)d - kd^2 - c^2 = cd - c^2$. Finally we have to add $r\delta$ the dimension of elementary transformations. The total dimension is:

$$(\text{****.}) \quad (r^2 - 1)(g - 1) + g + cd - c^2 - \delta.$$

Since $(\text{***}) \geq (\text{****})$ with equality iff $\delta = 0$, we can conclude surjectivity except for two problems. The first: what object have we calculated the dimension of? The second: what if $K^* \otimes L_i$ is special?

For the first problem we can form a universal Hom space, for maps from K to $W_k \overset{c}{\oplus} L_{k+1}$. The quotient comes for free. In general the Hom space is constructed as follows. Suppose we have families of bundles $\mathcal{E}$ over $S \times C$, and $\mathcal{F}$ over $T \times C$. Now consider $\mathcal{E}^* \otimes \mathcal{F}$ over $S \times T \times C$, and push down from C , then pull it back up and call it U . If $h^0(C, E^* \otimes F)$ is constant, then restricting to a point $(E \in S, F \in T)$ the bundle is $H^0(C, E^* \otimes F)$. And over a point $v \in H^0(C, E^* \otimes F)$ we get a morphism $E \rightarrow F$ over C . If $h^0(C, E^* \otimes F)$ is not constant we can take a stratification and check the dimension of each piece.

For the second problem we have $\deg(K^*) = \alpha$ with $1 \leq \alpha \leq 2g - 2$. So $g \geq 2$. We will do the cases $\alpha = 1$ and 2 separately. Then we will do the case $\alpha \geq 3$. For $\alpha = 1$, d must be 1 and $k \geq 2$. For fixed L_1 and L_2 distinct there are only finitely many line bundles K of $\deg = 0$ so that $h^0(C, K^* \otimes L_i) \neq 0$ for $i = 1$ and 2 . Furthermore we must have $d = 1$. The morphisms from K to L_i are unique up to automorphisms of W and hence there are only finitely many Q , one for each K . Now the dimension of elementary translations is r times the difference of degree or

$$r(r(g - 1) + d - 1) < \text{***}.$$

Now we assume $\alpha = 2$ and $h^0(C, K^* \otimes L_i) = 1$ for all i , and hence $d = 1$. The dimension of K is $\leq \alpha$. As above each K gives rise to a unique Q . So the dimension of Q is $\leq \alpha$ and adding the dimension of elementary transformations we get

$$\alpha + r(r(g - 1) + d - \alpha) < \text{***}.$$

Now we by reordering i if necessary, we can assume $h^0(C, K^* \otimes L_1) \geq 2$ and $\alpha = 2$. $h^0(C, K^* \otimes L_i) = 1$ for $i \neq 1$. We have two case. Case i) $d=2$ and hence $k=1$. Case ii) $d=1$. For case i, the dimension of K is zero. Furthermore, the dimension of Q is zero because there is only one fixed map from K to W up to automorphisms of W . Finally, the dimension of elementary transformations is

$$r(r(g-1) + d - 2) < ***.$$

For case ii, dimension K is again zero, but there is a one dimensional family of maps from K to L_1 up to automorphisms of W , and hence, there is a one dimensional family of Q . We add the dimension of elementary transformations to get

$$1 + r(r(g-1) + d - 2) < ***.$$

Now assume $3 \leq \alpha \leq 2g - 2$. By Clifford's Theorem, $h^0(C, K^* \otimes L_i) \leq \frac{\alpha}{2} + 1$. By Marten's Theorem the dimension of K with $h^0(C, K^* \otimes L_i) \leq \frac{\alpha}{2} + 1 - a$ equals $2a$ for a non-negative. Now assume all L_i have $h^0(C, K^* \otimes L_i) \leq \frac{\alpha}{2} + 1 - a$. The dimension of K is $2a$. The dimension of Q is $\leq kd(\frac{\alpha}{2} + 1 - a - d) + c(\frac{\alpha}{2} + 1 - a - c) + 2a$ or:

$$(kd + c)(\frac{\alpha}{2} + 1 - a - d) + cd - c^2 + 2a.$$

This is clearly maximal if $a = 0$, as $kd + c = r + 1 > 2$. Now adding the dimension of the elementary transformations and rearranging the terms we get:

$$r^2(g-1) + cd - c^2 + \frac{\alpha}{2}(r+1) - r\alpha - d + r + 1.$$

To show this is less than *** it suffices to show:

$$\frac{\alpha}{2}(r+1) - r\alpha - d + r + 1 < 1$$

$$\frac{-(r-1)}{2}\alpha - d + r + 1 < 1$$

$$-\frac{3}{2}r + \frac{3}{2} - d + r + 1 < 1$$

$$\frac{-r}{2} - d + \frac{5}{2} \leq \frac{1}{2} \quad \square$$

§3 THE MAIN THEOREM

Now we are prepared to prove the Main Theorem.

Proof of Main Theorem part A. Suppose $ed = r - 1$. By Proposition 1 and repeated use of Proposition 2, for a general $E_t \in SU$ we have a sequence:

$$0 \rightarrow W \rightarrow E_t \rightarrow A \rightarrow 0$$

for a fixed line bundle A . Now the family of E_t is given by elements of $\text{Ext}^1(W, A)$ modulo the automorphism of W . Ext^1 is given by $H^0(C, \omega_C \otimes W^* \otimes L)^*$. This decomposes into a direct sum of $H^0(C, \omega_C \otimes V_i^* \otimes A)^*$, which in turn is isomorphic to a direct sum of $H^0(C, \omega_C \otimes A \otimes L_i)^* \otimes U_i^*$. Concentrating on one i we can see that the quotient by the automorphisms of U_i gives a Grassmannian. So the whole thing is just a product of Grassmannians.

Now for $ed = r + 1$, Proposition 3 shows that for a general $E_t \in SU$, we have an exact sequence

$$0 \rightarrow A \rightarrow W \rightarrow E_t \rightarrow 0,$$

for some fixed bundle A . So the family is parameterized by $\text{Hom}(A, W)$ modulo the automorphisms on W . We can decompose and get a direct sum of $H^0(C, L^* \otimes L_i) \otimes U_i$

modulo the automorphisms of W — but again, this is just a product of Grassmannians. And we are done. $\square$

To prove part B we need a result of Boden and Yokogawa.

Definition. A variety X is stably rational of level k if $X \times \mathbb{P}^k$ is rational.

Theorem 3 (Boden and Yokogawa [3]). *If $(r, d) = 1$ then $SU(r, d)$ is stably rational of level $r - 1$*

Now we prove part B.

Proof of Main Theorem part B. As in the argument for part A, using Propositions 1 and 2 repeatedly we get a sequence for general E_t ,

$$0 \rightarrow W \rightarrow E_t \rightarrow F_u \rightarrow 0,$$

where F_u is stable. This means a general E_t is an extension of a stable bundle F_u by W . So for a general F_u and a general extension by W we get $E_t \in SU$. The map to SU is not one to one, but its fiber is just the automorphisms of W since for general E_t , W is a unique subbundle.

So now we need to construct the map more carefully, then take care of the fiber. First note that by the hypothesis the rank and degree of F_u (which is $(\delta, r(g-1)+d)$) are coprime because all factors of δ divide r and not d . So now let $SU' = SU(\delta, r(g-1)+d)$. SU' is a fine moduli space with a Poincare bundle $\mathcal{P}$ over $SU' \times C$. We also have projections p_1 and p_2 onto SU' and C respectively. Now we will form a bundle parameterizing extensions of $F_u \in SU'$ by W . Consider

$$p_2^* p_{2*} (p_1^* (W^* \otimes \omega_C) \otimes \mathcal{P})^*.$$

The general fiber is $H^0(C, W^* \otimes \omega_C \otimes F_u)^*$ which is canonically $\text{Ext}^1(F_u, W)$. It is also canonically $\bigoplus^e H^0(C, L_i^* \otimes \omega_C \otimes F_u)^* \otimes U_i^*$. By taking an affine open subset on SU' , we can

assume the bundle is trivial, and that we have a universal extension space parameterized by the total space of the bundle, which by restricting to a smaller affine open subspace of SU' is trivial and hence has no higher cohomology which satisfies the conditions of [Lemma 2.4, 7]. And therefore this gives a dominant map from the bundle over SU' to SU which has fiber isomorphic to $\text{Aut}(W)$. But clearly the action of $\text{Aut}(W)$ on the bundle over SU gives a product of Grassmannians cross SU' . The dimension of the product of Grassmannians is the rank of the bundle minus ed^2 . For the rank of the bundle we have

$$h^0(\omega_C \otimes L_i^* \otimes F_u) = \delta(g-1) + (r-\delta)(g-1) + d + 2(g-1)\delta - \delta(g-1).$$

This is $(r+\delta)(g-1)+d$. Now to get the rank of the full bundle we multiply by e the number of L_i , and d the rank of U_i . But $ed = r-\delta$, so we get $(r+\delta)(r-\delta)(g-1)+ed^2$. Now $\text{Aut}(W)$ has dimension ed^2 . So the product of Grassmannians has dimension $(r+\delta)(r-\delta)(g-1) \geq r-1$. Now by Boden Yokogawa SU' cross the product of Grassmannians is rational. And therefore so is SU . $\square$

Remark 3. The condition that $ed = r - \delta$ where all prime divisors of δ are divisors of r is necessary. Suppose to the contrary that a prime p divides δ but p does not divide r , then we can solve the equation $r(g-1) + d \equiv 0 \pmod{p}$. The rank and degree would then not be relatively prime for those g which solve the equation.

Remark 4. Solutions for g that satisfy the above remark are the only g where rationality can fail, because otherwise $(r(g-1) + d, \delta) = 1$ and the proof goes through. For a given g , it is enough if $(r(g-1) + d, \delta) = 1$ for one possible δ . In the case of $r = 13$ and $d = 5$ or $d = 8 = (13 - 5)$ (the first case where $(r, d) = 1$ but the hypothesis of the Main Theorem does not hold) we get three equations. First the Boden Yokogawa equations $(g, 5) = 1$ and $(g, 8) = 1$. The Chinese Remainder Theorem gives $g \equiv 0 \pmod{10}$. Now we let $e = 2$ so $\delta = 13 - 2(5) = 3$ and we get $13(g-1) + 5 \equiv 0 \pmod{3}$. The final equation is then

$$g \equiv 20 \pmod{30}.$$

We shall have more to say about this in the next section, after making a conjecture.

§4 CONJECTURES

Inspired by the Main Theorem, it seems natural to prove that SU is rational if $ed = r + \delta$ where δ has the property that any prime divisor of δ is a prime divisor of r , and $0 < d < r$.

We **attempt** a proof.

We begin by constructing the universal hom space over SU and $\{W\}$. It might have more than one irreducible component, corresponding to bundles E_t where $h^0(C, W^* \otimes E_t)$ is large because h^1 is large. But for a general bundle we have the above $h^1 = 0$. So we get one irreducible component which dominates SU . Now given a morphism $W \rightarrow E$. We get a morphism $K_{E,W} \hookrightarrow W$. We would like $K_{E,W}$ to be stable. And given $SU' = SU(\delta, -r(g-1) - d)$, there is a dominant map from SU to SU' . This would imply that $\delta < r$ since if it is bigger than r the map SU to SU' cannot dominate for dimension reasons. And if $\delta = r$ we get birationally isomorphic spaces SU and SU' .

But how do we know $K_{E,W}$ is stable for a general E when $0 < \delta < r$? The problem is we do not! So for now, **suppose** that $K = K_{E,W}$ is stable. For a general K we have $h^1(C, K^* \otimes W) = 0$. So we calculate the dimension of the irreducible component of the universal Hom space containing $\{K | h^1(C, K^* \otimes W) = 0\}$. First we count the dimension of $\text{Hom}(K, W)$.

$$\begin{aligned} h^0(C, K^* \otimes W) &= \delta ed \left(\frac{r(g-1) + d}{\delta} - (g-1) \right) \\ &= ed(r - \delta)(g-1) + ed^2 \\ &= (r^2 - \delta^2)(g-1) + ed^2 \end{aligned}$$

Adding $\dim U' = \dim U(\delta, -r(g-1) - d) = (\delta^2(g-1))$, we get $r^2(g-1) + ed^2$. This is the dimension of U plus the dimension of $\text{Aut } W$. So we get a dominant map from the

irreducible component of the Universal Hom space dominating SU' to SU and the general fiber is $\text{Aut}(W)$.

Since $(\delta, -r(g-1)-d) = 1$ because all prime divisors of δ divide r , we can then restrict to an open set of SU' and construct the Universal Hom bundle by Taking the pullback of the pushdown of the Poincare bundle $\mathcal{P}$ tensored by the pull back of W^* . This bundle is locally trivial and has fibers $H^0(C, E \otimes W^*)$ which equals the direct sum $\bigoplus^e H^0(C, E \otimes L_i^*) \otimes U_i$. Now the action of $\text{Aut}(W)$ gives SU' cross a product of Grassmannians. Now SU is rational because of Boden Yokogawa and the fact that the dimension of the product of Grassmannians is $\geq r - 1$.

Conjecture 1. *Let C be a smooth projective irreducible curve over $\mathbb{C}$. If $(r, d) = 1$ and d divides $r + \delta$ where $0 < \delta < r$ and every prime divisor of δ divides r , then $SU = SU(r, r(g-1) + d)$ is rational.*

As noted in the above plausibility argument, Conjecture 1 would follow from the stability of the bundle $K_{E,W}$ and the vanishing of $h^1(C, K_{E,W}^* \otimes W)$

But it is actually enough to show the vanishing of one $h^1(C, K_{E,W}^* \otimes W)$ provided $h^1(C, W^* \otimes E) = 0$. This is because we would then get a family of $K_{E,W}$ for which $h^1(C, K_{E,W}^* \otimes W) = 0$ generically. The dimension of that component of the Universal Hom is, by the calculations preceding Conjecture 1, the dimension of the family $+(r^2 - \delta^2)(g-1) + ed^2$. Now this adds up to $r^2(g-1) + 1 + ed^2$ (the dimension of $\text{Hom}(W, E)$) if and only if the dimension of the family is $\delta^2(g-1) + 1$. But $\deg K_{W,E}$ is fixed, and $\mu^{\max} < 0$ because it is a subbundle of W . This means there are only finitely many HN types. But the dimension of a parameter space for unstable bundles is less than $\delta^2(g-1) + 1$. Hence, $K_{W,E}$ is generically stable. So it would suffice to prove the following conjecture in order to prove Conjecture 1.

Conjecture 2. *Let C be as above. There is an $E \in SU$ such that $h^1(C, W^* \otimes E) = 0$, E*

is spanned by W , and $h^1(C, K_{W,E}^* \otimes W) = 0$.

We can prove Conjecture 2 if $e = 1$ or 2 .

Proposition 4. *Conjecture 2 holds if $e = 1$ or 2 .*

To prove this we need a Lemma belonging to folklore. By an elementary transformation of a bundle E we mean the kernel, E_p , of a sequence.

$$0 \rightarrow E_p \rightarrow E \rightarrow \mathcal{O}_p \rightarrow 0,$$

where p is a reduced point on C . This corresponds to blowing up a point $p \in \mathbb{P}(E)$ and taking the pushdown of the fiber. Corresponding to a point on $\mathbb{P}(E)$ there is a $r - 2$ plane in $\mathbb{P}(E_p)$ and this corresponds to a point in $\mathbb{P}(E_p^*)$. If we blow up this point, blow down the fiber, and then dualize we get E back. This is known as an inverse elementary transformation.

Lemma 1. *Let C be as above. For a general bundle $E \in U(r, d)$ or $E \in SU(r, d)$ a general elementary transformation is stable and so is a general inverse elementary transformation.*

Proof of Lemma 1. Suppose to the contrary that the general E_p and hence all E_p are unstable. E_p has fixed degree $(d - 1)$ and $\mu^{\max}(E_p) < \frac{d}{r}$ because E_p is a subbundle of the stable bundle E . This means there are finitely many HN types.

Now consider the parameter space of a general E_p with its universal bundle $\mathcal{E}$. Take $X = \mathbb{P}(\mathcal{E})$. A point on X is an elementary transformation of a bundle F^* which when dualized gives a bundle $E \in U(r, d)$. If the map to $U(r, d)$ does not dominate then the Lemma is proved. So assume the map dominates. The dimension of X is the dimension of the parameter space $+r - 1$. Which equals (by Sundaram) $r^2(g - 1) - r + 1$ for the parameter space $+r - 1 = r^2(g - 1)$. That means the fiber for a general bundle $E \in U(r, d)$ consists of finitely many points. But those points represent all the elementary transformations of E which are not stable (because every such transformation has an

inverse). The conclusion is a general elementary transformation of a stable bundle is stable. This proves the Lemma. $\square$

Proof of Proposition 4. In the case $e = 1$ there is only one linebundle L_1 . Twisting by L_1^* we can assume the bundle is $\mathcal{O}_C$ and W is $H^0(C, E) \otimes \mathcal{O}_C$. We will now construct a bundle for $d = 2r$ then work our way down in degree. Let A be a general line bundle of $\deg g + 1$. A is non special and spanned. So let E be the direct sum of r copies of A . $K_{E,W}^* = E$ and so it is nonspecial. The only problem is that the original bundle E is not stable. We can choose a one parameter family of stable bundles E_t degenerating to $E_0 = E$. A general E_t will be stable and spanned and so will have a bundle $K_{E_t,W}$. So taking a possibly smaller family we parameterize a family of $K_{E_t,W}^*$ with a general bundle nonspecial. This finishes the case $d = 2r$.

To get the case $d = 2r - 1$ we use elementary transformations. We can assume that a general $E_t \in M_r = U(r, rg + r)$ has $K_{E,W}^*$ non special. Furthermore, a general elementary transform of E_t is stable by Lemma 1. And for $F \in M_{r-1} = U(r, rg + r - 1)$, a general inverse transformation is stable by Lemma 1, and a general bundle F is spanned. Putting these together shows that there is a pair $E \in M_r$ and $E_p \in M_{r-1}$. Where E is spanned with $K_{E,W}^*$ nonspecial and E_p an elementary transformation of E with E_p spanned. We have an exact sequence

$$0 \rightarrow E_p \rightarrow E \rightarrow \mathcal{O}_p \rightarrow 0.$$

There is a second exact sequence

$$0 \rightarrow H^0(C, E_p) \otimes \mathcal{O}_C \rightarrow H^0(C, E) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C \rightarrow 0.$$

The second sequence surjects onto the first and the kernel is

$$0 \rightarrow K_{E_p, W_p} \rightarrow K_{E, W} \rightarrow \mathcal{O}_C(-p) \rightarrow 0.$$

W_p is just $H^0(C, E_p) \otimes \mathcal{O}_C$. And by dualizing the last sequence and taking cohomology we see that K_{E_p, W_p}^* is nonspecial. Repeating this argument proves the case $e = 1$.

For the case $e = 2$ there are line bundles L_1 and L_2 of deg 0. Now do the case where $d = r$. Choose a general line bundle A with $\deg(A) = g$ so that $A \otimes L_i^*$ is not special for $i = 1$ or 2 . And choose A such that the divisors $|A \otimes L_1^*|$ and $|A \otimes L_2^*|$ have no points in common. Now let E be the direct sum of r copies of A . It is easily verified that $h^1(C, K_{E,W}^* \otimes L_i) = 0$ for $i = 1$ or 2 . This proves the case $d = r$. The remaining cases are proved as above using elementary transformations. $\square$

Remark 5. The case $e = 1$ has no impact on Conjecture 1. The case $e = 2$ seems to but does not. For suppose that $2d = r + \delta$ where every prime factor of δ divides r . Then $2(r - d) = r - \delta$ where δ has the above properties. So the result is already known because $SU(r, r(g - 1) + d) \cong SU(r, r(g - 1) + r - d)$.

However the case $e = 1$ may have some use. Throughout this paper we assumed the choice of bundles $\{L_1, \dots, L_e\}$ was of arbitrary distinct bundles. But we never needed that fact (outside of distinctness). So consider L_1 such that $L_1^{\otimes i}$ is trivial iff i is a multiple of e . Now set $L_i = L_1^{\otimes i}$. The set of L_i corresponds to a cyclic étale cover $\tilde{C}$ of C . The L_i pull back to be trivial, so $\tilde{W}$ the pullback of W is trivial. On $\tilde{C}$, a general $K_{\tilde{E}, \tilde{W}}^*$ is nonspecial. So we expect (but have not proven!) that for general $\tilde{E}$ a pullback of E on C , $K_{\tilde{E}, \tilde{W}}^*$ is nonspecial. Proving this would prove Conjecture 2 and hence Conjecture 1. Because then $K_{E,W} \otimes L_i$ would be nonspecial for all i .

To prove this last conjecture it is required to prove that a general $\tilde{E}$ is spanned. That much we can do. But unfortunately we work over C and not $\tilde{C}$. A new proof of the Second Theorem may lead to a proof of the above.

Now we prove the Second Theorem.

Proof of Second Theorem. A cyclic cover of degree e (where p does not divide e) corresponds to a subgroup of e torsion points of the Jacobian. Or in other words a set:

$$\{L, L^{\otimes 2} \dots L^{\otimes (e-1)}, L^{\otimes e}\},$$

with $L^{\otimes i} = \mathcal{O}_C$ iff $i = e$. Setting the above set of line bundles equal to $L_1, L_2, \dots, L_e$, and noting that the sections of π^*E are the pullback of W , we then get the Second Theorem from Proposition 3. $\square$

Remark 6. Assuming the conjecture on the stability of $K_{E,W}$ there are more equations (as in Remark 4) that g must solve. We have $r = 13$ and $d = 5$ or 8 . The case $e = 2$ gives nothing new (as in Remark 5). But for $e = 3$ we have $3 \times 8 = 24 = 13 + 11$ which gives the equation $13(g - 1) + 8 \equiv 0 \pmod{11}$ or

$$g \equiv 8 \pmod{11}.$$

But $e = 3$ and $d = 5$ gives nothing new. However, for $e = 4$ and $d = 5$ we get $20 = 13 + 7$ so $13(g - 1) + 5 \equiv 0 \pmod{7}$. This reduces to

$$g \equiv 6 \pmod{7}.$$

Taking these equations, the previous equations, and using the Chinese Remainder Theorem we get

$$g \equiv 2, 120 \pmod{2, 310}.$$

REFERENCES

1. Ballico, *Stable rationality for the variety of vector bundles over an algebraic curve*, J. Lond. Math. Soc. (2) **30** (1984), 21 – 26.
2. A. Beauville, J-L. Colliot-Thélène, J-J Sansuc and P. Swinnerton-Dyer, *Variétés stablement rationnelles non rationnelles*, Ann. of Mathematics **121** (1985), 283 – 318.
3. H.U. Boden and K. Yokogawa, *Rationality of moduli spaces of parabolic bundles*, Jour. of Lond. Math. Soc. (to appear).
4. H. Lange, *Universal families of extensions*, Jour. of Alg. **83**, 101 – 112.
5. M.S. Narasimhan and C.S. Seshadri, *Stable and unitary vector bundles on a compact Riemann surface*, Ann. of Math. **82** (1969), 540 – 567.
6. P.E. Newstead, *Rationality of moduli spaces of stable bundles*, Math. Ann. **215** (1975), 251 – 268; *Correction*, Math. Ann. **249** (1980), 281 – 282.
7. S. Ramanan, *The moduli spaces of vector bundles over an algebraic curve*, Math. Ann. **200** (1973), 69 – 84.
8. N. Sundaram, *Special divisors and vector bundles*, Tohuko Math. J. **39** (1987), 175 – 223.
9. A.N. Tyurin, *On the classification of two dimensional vector bundles over an algebraic curve of arbitrary genus*, Izv. Akad. Nauk. SSSR Ser. Mat. **28** (1964), 21 – 52.

10. ———, *Classification of vector bundles over an algebraic curve of arbitrary genus*, Izv. Akad. Nauk. SSSR Ser. Mat. **29** (1964), 657 – 688; *English translation*, Amer. Math. Soc. Trans. **63** (1967), 245 – 279.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD,
BOMBAY, INDIA

E-MAIL: BUTLER@MATH.TIFR.RES.IN